

66/02/2T
 12/20/99
 J674 U.S. PTO

U.S. PTO
 09/466935
 12/20/99

UTILITY PATENT APPLICATION TRANSMITTAL <small>(Only for new nonprovisional applications under 37 CFR 1.53(b))</small>	Attorney Docket No.	0010-1070-0
	First Inventor or Application Identifier	Vitaliy A. LIVSHITS, et al.
	Title	NOVEL GENE AND METHOD FOR PRODUCING L-AMINO ACIDS

APPLICATION ELEMENTS <small>See MPEP chapter 600 concerning utility patent application contents</small>	ADDRESS TO: Assistant Commissioner for Patents Box Patent Application Washington, DC 20231
1. <input checked="" type="checkbox"/> Fee Transmittal Form (e.g. PTO/SB/17) <small>(Submit an original and a duplicate for fee processing)</small> 2. <input checked="" type="checkbox"/> Specification Total Pages 45 3. <input checked="" type="checkbox"/> Drawing(s) (35 U.S.C. 113) Total Sheets 4 4. <input checked="" type="checkbox"/> Oath or Declaration Total Pages 3 a. <input checked="" type="checkbox"/> Newly executed (original) b. <input type="checkbox"/> Copy from a prior application (37 C.F.R. §1.63(d)) <small>(for continuation/divisional with box 15 completed)</small> i. <input type="checkbox"/> DELETION OF INVENTOR(S) Signed statement attached deleting inventor(s) named in the prior application, see 37 C.F.R. §1.63(d)(2) and § 1.33(b). 5. <input type="checkbox"/> Incorporation By Reference <small>(usable if box 4B is checked)</small> The entire disclosure of the prior application, from which a copy of the oath or declaration is supplied under Box 4B, is considered to be part of the disclosure of the accompanying application and is hereby incorporated by reference therein.	ACCOMPANYING APPLICATION PARTS 6. <input type="checkbox"/> Assignment Papers (cover sheet & document(s)) 7. <input type="checkbox"/> 37 C.F.R. §3.73(b) Statement <input type="checkbox"/> Power of Attorney <small>(when there is an assignee)</small> 8. <input type="checkbox"/> English Translation Document <small>(if applicable)</small> 9. <input type="checkbox"/> Information Disclosure Statement (IDS)/PTO-1449 <input type="checkbox"/> Copies of IDS Citations 10. <input type="checkbox"/> Preliminary Amendment 11. <input checked="" type="checkbox"/> White Advance Serial No. Postcard 12. <input type="checkbox"/> Small Entity Statement(s) <input type="checkbox"/> Statement filed in prior application. Status still proper and desired. 13. <input checked="" type="checkbox"/> Certified Copy of Priority Document(s)(1) <small>(if foreign priority is claimed)</small> Notice of Priority, Receipt of an Original Deposit of Microorganisms for the Purposes of Patent Procedure (VKPM B-7680, VKPM B-7681, VKPM B-7682, VKPM B-7683, VKPM B-7684, VKPM B-7685, VKPM B-7700) 14. <input checked="" type="checkbox"/> Other:
15. If a CONTINUING APPLICATION, check appropriate box, and supply the requisite information below: <input type="checkbox"/> Continuation <input type="checkbox"/> Divisional <input type="checkbox"/> Continuation-in-part (CIP) of prior application no.: Prior application information: Examiner: Group Art Unit:	
16. Amend the specification by inserting before the first line the sentence: <input type="checkbox"/> This application is a <input type="checkbox"/> Continuation <input type="checkbox"/> Division <input type="checkbox"/> Continuation-in-part (CIP) of application Serial No. Filed on <input type="checkbox"/> This application claims priority of provisional application Serial No. Filed	
17. CORRESPONDENCE ADDRESS OBLON, SPIVAK, MCCLELLAND, MAIER & NEUSTADT, P.C. FOURTH FLOOR 1755 JEFFERSON DAVIS HIGHWAY ARLINGTON, VIRGINIA 22202 (703) 413-3000 FACSIMILE: (703) 413-2220	

Name:	Norman F. Oblon	Registration No.:	24,618
Signature:	MARVIN J. SPIVAK REGISTRATION NUMBER 24,913	Date:	12/20/99
Name:		Registration No.:	

NOVEL GENE AND METHOD FOR PRODUCING L-AMINO ACIDS

Technical field

The present invention relates to biotechnology, and
5 more specifically to a method for producing amino acid,
especially for a method for producing L-homoserine, L-
threonine, L-valine or L-leucine using a bacterium
belonging to the genus *Escherichia*.

10 Background Art.

The present inventors obtained, with respect to *E.*
coli K-12, a mutant having mutation, *thrR* (herein
referred to as *rhtA23*) that is concerned in resistance
to high concentrations of threonine or homoserine in a
15 minimal medium (Astaurova, O.B. et al., Appl. Bioch. And
Microbiol., 21, 611-616 (1985)). The mutation improved
the production of L-threonine (SU Patent No. 974817),
homoserine and glutamate (Astaurova, O.B. et al., Appl.
Bioch. And Microbiol., 27, 556-561, 1991) by the
20 respective *E. coli* producing strains.

Furthermore, the present inventors has revealed that
the *rhtA* gene exists at 18 min on *E. coli* chromosome and
that the *rhtA* gene is identical to ORF1 between *pexB* and
ompX genes. The unit expressing a protein encoded by the
25 ORF has been designated as *rhtA* (*rht*: resistance to

homoserine and threonine) gene. The *rhtA* gene includes a 5'-noncoding region including SD sequence, ORF1 and a terminator. Also, the present inventors have found that a wild type *rhtA* gene participates in resistance to threonine and homoserine if cloned in a multicopy state and that the *rhtA23* mutation is an A-for-G substitution at position -1 with respect to the ATG start codon (ABSTRACTS of 17th International Congress of Biochemistry and Molecular Biology in conjugation with 1997 Annual Meeting of the American Society for Biochemistry and Molecular Biology, San Francisco, California August 24-29, 1997, abstract No. 457).

It is found that at least two different genes which impart threonine and homoserine resistance in a multicopy state exist in *E. coli* during cloning of the *rhtA* gene. One of the genes is the *rhtA* gene, and the other gene was found to be *rhtB* gene which confers homoserine resistance (Russian Patent Application No. 98118425).

20

Disclosure of the Invention

An object of the present invention is to provide a method for producing an amino acid, especially, L-homoserine, L-threonine and branched chain amino acids

25

with a higher yield.

The inventors have found that a region at 86 min on *E. coli* chromosome, when cloned by a multicopy vector, impart resistance to L-homoserine to cells of *E. coli*.

5 the inventors further found that there exists in the upstream region another gene, *rhtC*, which involves resistance to threonine, and that when these genes are amplified, the amino acid productivity of *E. coli* can be improved like the *rhtA* gene. On the basis of these
10 findings, the present invention have completed.

Thus, the present invention provides:

(1) A bacterium belonging to the genus *Escherihia*, wherein L-threonine resistance of the bacterium is enhanced by enhancing an activity of
15 protein as defined in the following (A) or (B) in a cell of the bacterium:

(A) a protein which comprises the amino acid sequence of SEQ ID NO: 4; or

(B) a protein which comprises the amino acid
20 sequence including deletion, substitution, insertion or addition of one or several amino acids in the amino acid sequence of SEQ ID NO: 4, and which has an activity of making a bacterium having the protein L-threonine-resistant;

25 (2) The bacterium according to (1), wherein L-

homoserine resistance of the bacterium is further enhanced by enhancing an activity of protein as defined in the following (C) or (D) in a cell of the bacterium:

(C) a protein which comprises the amino acid
5 sequence of SEQ ID NO: 2; or

(D) a protein which comprises the amino acid
sequence including deletion, substitution, insertion or
addition of one or several amino acids in the amino acid
sequence of SEQ ID NO: 2, and which has an activity of
10 making a bacterium having the protein L-homoserine-
resistant;

(3) The bacterium according to (1) or (2), wherein
the activity of protein as defined in (A) or (B) is
enhanced by transformation of the bacterium with DNA
15 coding for the protein as defined in (A) or (B);

(4) The bacterium according to (2), wherein the
activity of protein as defined in (C) or (D) is enhanced
by transformation of the bacterium with DNA coding for
the protein as defined in (C) or (D);

20 (5) A method for producing an amino acid, comprising
the steps of:

cultivating the bacterium as defined in any one of
(1) to (4), which has an ability to produce an amino
acid, in a culture medium, to produce and accumulate the
25 amino acid in the medium, and recovering the amino acid

from the medium;

(6) The method according to (5), wherein the amino acid is selected from the group consisting of L-homoserine, L-threonine and branched chain amino acids;

5 (7) The method according to (6), the branched chain amino acid is L-valine or L-leucine.

(8) A DNA which encode a protein defined in the following (A) or (B):

(A) a protein which has the amino acid sequence of
10 SEQ ID NO: 4;

(B) a protein which has the amino acid sequence of
SEQ ID NO: 4 including substitution, deletion, insertion,
addition, or inversion of one or several amino acids,
and has an activity of making a bacterium having the
15 protein L-threonine-resistant.

9. The DNA of (8) which is a DNA defined in the following (a) or (b):

(a) a DNA which comprises the nucleotide sequence of nucleotide numbers 187 to 804 in SEQ ID NO: 3;

20 (b) a DNA which is hybridizable with a nucleotide sequence of nucleotide numbers 187 to 804 in SEQ ID NO: 3 or a probe prepared from the nucleotide sequence under a stringent condition, and encodes a protein having an activity of making a bacterium having the protein L-
25 threonine-resistant; and

10. The DNA of (9) wherein the stringent condition is a condition in which washing is performed at 60 °C, and at a salt concentration corresponding to 1 x SSC and 0.1 % SDS.

5 The DNA fragment coding for the protein as defined in the above (A) or (B) may be referred to as "*rhtC* gene", a protein coded by the *rhtC* gene may be referred to as "*RhtC* protein", the DNA coding for the protein as defined in the above (C) or (D) may be referred to
10 as "*rhtB* gene", a protein coded by the *rhtB* gene may be referred to as "*RhtB* protein". An activity of the *RhtC* protein which participate in resistance to L-threonine of a bacterium (i.e. an activity of marking a bacterium having the *RhtC* protein L-threonine-resistant) may be
15 referred to as "Rt activity", and an activity of the *RhtB* protein which participates in resistance to L-homoserine of a bacterium (i.e. an activity of marking a bacterium having the *RhtB* protein L-homoserine-resistant) may be referred to as "Rh activity". A
20 structural gene encoding the *RhtC* protein or *RhtB* protein in the *rhtC* gene or *rhtB* gene may be referred to as "*rhtC* structural gene" or "*rhtB* structural gene". The term "enhancing the Rt activity or the Rh activity" means imparting resistance to threonine or homoserine to
25 a bacterium or enhance the resistance by means of

increasing the number of molecules of the RhtC protein or RhtB protein increasing a specific activity of these proteins, or desensitizing negative regulation against the expression or the activity of these proteins or the like. The terms "DNA coding for a protein" mean a DNA of which one of strands codes for the protein when the DNA is double-stranded. The L-threonine resistance means a property that a bacterium grows on a minimal medium containing L-threonine at a concentration at which a wild-type strain thereof not grow, usually at >30 mg/ml. The L-homoserine resistance means a property that a bacterium grows on a minimal medium containing L-homoserine at a concentration at which a wild-type strain thereof not grow, usually at >5 mg/ml. The ability to produce an amino acid means a property that a bacterium produce and accumulates the amino acid in a medium in a larger amount than a wild type strain thereof.

According to the present invention, resistance to threonine, or threonine and homoserine of a high concentration can be imparted to a bacterium belonging to the genus *Escherichia*. A bacterium belonging to the genus *Escherichia*, which has increasing resistance to threonine, or threonine and homoserine and an ability to accumulate an amino acid, especially, L-homoserine, L-

threonine, or branched chain amino acids such as L-valin and L-leucine in a medium with a high yield.

The present invention will be explained in detail below.

5 <1> DNA used for the present invention

The first DNA fragment used for the present invention (*rhtC* gene) coding for a protein having the Rt activity and having the amino acid sequence of SEQ ID NO: 4. Specifically, the DNA may be exemplified by a DNA
10 comprising a nucleotide sequence of the nucleotide numbers 187 to 804 of a nucleotide sequence of SEQ ID NO: 3.

The second DNA fragment used for the present invention (*rhtB* gene) coding for a protein having the Rh
15 activity an having the amino acid sequence of SEQ ID NO: 2. Specifically, the DNA may be exemplified by a DNA comprising a nucleotide sequence of the nucleotide numbers 557 to 1171 of a nucleotide sequence of SEQ ID NO: 1.

20 The *rhtB* gene having the nucleotide sequence of SEQ ID NO: 1 corresponds to a part of sequence complement to the sequence of GenBank accession number M87049, and includes fl38 (nucleotide numbers 61959-61543 of M87049) which is a known but function-unknown ORF (open reading
25 frame) present at 86 min on *E. coli* chromosome, and 5'-

and 3'-flanking regions thereof. The f138, which had only 160 nucleotides in the 5'-flanking region, could not impart the resistance to homoserine. No termination codon is present between the 62160 and 61959 nucleotides of M87049 (upstream the ORF f138). Moreover, one of the ATG codons of this sequence is preceded by a ribosome-binding site (62171-62166 in M87049). Hence, the coding region is 201 bp longer. The larger ORF (nucleotide numbers 62160 to 61546 of M87049) is designated as *rhtB* gene.

The *rhtB* gene may be obtained, for example, by infecting Mucts lysogenic strain of *E. coli* using a lysate of a lysogenic strain of *E. coli* such as K12 or W3110 according to the method in which mini-Mu d5005 phagemid is used (Groisman, E.A., et al., J. Bacteriol., 168, 357-364 (1986)), and isolating phagemid DNAs from colonies growing on a minimal medium containing kanamycin (40 µg/ml) and L-homoserine (10 mg/ml). As illustrated in the Example described below, the *rhtB* gene was mapped at 86 min on the chromosome of *E. coli*. Therefore, the DNA fragment including the *rhtB* gene may be obtained from the chromosome of *E. coli* by colony hybridization or PCR (polymerase chain reaction, refer to White, T.J. et al, Trends Genet. 5, 185 (1989)) using oligonucleotide(s) which has a sequence corresponding to

the region near the portion of 86 min on the chromosome *E. coli*.

Alternatively, the oligonucleotide may be designed according to the nucleotide sequence of SEQ ID NO: 1. By
5 using oligonucleotides having nucleotide sequences corresponding to an upstream region from the nucleotide number 557 and a downstream region from the nucleotide number 1171 in SEQ ID NO: 1 as the primers for PCR, the entire coding region can be amplified.

10 Synthesis of the oligonucleotides can be performed by an ordinary method such as a phosphoramidite method (see *Tetrahedron Letters*, 22, 1859 (1981)) by using a commercially available DNA synthesizer (for example, DNA Synthesizer Model 380B produced by Applied Biosystems).
15 Further, the PCR can be performed by using a commercially available PCR apparatus (for example, DNA Thermal Cycler Model PJ2000 produced by Takara Shuzo Co., Ltd.) using *Taq* DNA polymerase (supplied by Takara Shuzo Co., Ltd.) in accordance with a method designate by the
20 supplier.

The *rhtC* gene was obtained in the DNA fragment including *rhtB* gene by chance when *rhtB* was cloned as described later in the embodiments. The *rhtC* gene corresponds to a corrected, as described below, sequence
25 of O128 (nucleotide numbers 60860-61480 of GeneBank

accession number M87049) which is a known but function-unknown ORF. The *rhtC* gene may be obtained by PCR or hybridization using oligonucleotides designed according to the nucleotide sequence of SEQ ID NO: 3. By using
5 oligonucleotides having nucleotide sequence corresponding to a upstream region from nucleotide number 187 and a downstream region from the nucleotide number 804 in SEQ ID NO: 3 as the primers for PCR, the entire coding region can be amplified.

10 In the present invention, the DNA coding for the RhtB protein of the present invention may code for RhtB protein including deletion, substitution, insertion, or addition of one or several amino acids at one or a plurality of positions, provided that the Rh activity of
15 RhtB protein encoded thereby is not deteriorated. Similarly, the DNA coding for the RhtC protein of the present invention may code for RhtC protein including deletion, substitution, insertion, or addition of one or several amino acids at one or a plurality of positions,
20 provided that the Rt activity of RhtC protein encoded thereby is not deteriorated.

The DNA, which codes for the substantially same protein as the RhtB protein or RhtC protein as described above, may be obtained, for example, by modifying the
25 nucleotide sequence, for example, by means of the site-

directed mutagenesis method so that one or more amino acid residues at a specified site involve deletion, substitution, insertion, or addition. DNA modified as described above may be obtained by the conventionally known mutation treatment. The mutation treatment includes a method for treating a DNA coding for the RhtB protein or RhtC protein *in vitro*, for example, with hydroxylamine, and a method for treating a microorganism, for example, a bacterium, belonging to the genus *Escherichia* harboring a DNA coding for the RhtB protein with ultraviolet irradiation or a mutating agent such as N-methyl-N'-nitro-N-nitrosoguanidine (NTG) and nitrous acid usually used for the mutation treatment.

The DNA, which codes for substantially the same protein as the RhtB protein or RhtC protein, can be obtained by expressing a DNA subjected to *in vitro* mutation treatment as described above in multicopy in an appropriate cell, investigating the resistance to homoserine or threonine, and selecting the DNA which increase the resistance.

It is generally known that an amino acid sequence of a protein and a nucleotide sequence coding for it may be slightly different between species, strains, mutants or variants.

Therefore the DNA, which codes for substantially the

same protein as the RhtC protein, can be obtained by isolating a DNA which hybridizes with DNA having, for example, a nucleotide sequence of the nucleotide numbers 187 to 804 of the nucleotide sequence of SEQ ID NO: 3 or
5 a probe obtainable therefrom under stringent conditions, and which codes for a protein having the Rt activity from a bacterium belonging to the genus *Escherihia* which is subjected to mutation treatment, or a spontaneous mutant or a variant of a bacterium belonging to the
10 genus *Escherihia*.

Also, the DNA, which codes for substantially the same protein as the RhtB protein, can be obtained by isolating a DNA which hybridizes with DNA having, for example, a nucleotide sequence of the nucleotide numbers
15 557 to 1171 of the nucleotide sequence of SEQ ID NO: 1 or a probe obtainable therefrom under stringent conditions, and which codes for a protein having the Rh activity, from a bacterium belonging to the genus *Escherichia* which is subjected to mutation treatment, or
20 a spontaneous mutant or a variant of a bacterium belonging to the genus *Escherichia*.

The term "stringent conditions" referred to herein is a condition under which so-called specific hybrid is formed, and non-specific hybrid is not formed. It is
25 difficult to clearly express this condition by using any

numerical value. However, for example, the stringent conditions include a condition under which DNAs having high homology, for example, DNAs having homology of not less than 70% with each other are hybridized, and DNAs
5 having homology lower than the above with each other are not hybridized. Alternatively, the stringent condition is exemplified by a condition under which DNA's are hybridized with each other at a salt concentration corresponding to an ordinary condition of washing in
10 Southern hybridization, i.e., 60 °C, 1 x SSC, 0.1 % SDS, preferably 0.1 x SSC, 0.1 % SDS.

<2> Bacterium belonging to the genus *Escherichia* of the present invention

15 The bacterium belonging the genus *Escherichia* of the present invention is a bacterium belonging to the genus *Escherichia* of which the Rt activity is enhanced. Preferred embodiment of the bacterium of the present invention is a bacterium which is further enhanced the
20 Rh activity. A bacterium belonging to the genus *Escherichia* is exemplified by *Escherichia coli*. The Rt activity can be enhanced by, for example, amplification of the copy number of the *rhtC* structural gene in a cell, or transformation of a bacterium belonging to the genus
25 *Escherihia* with a recombinant DNA in which a DNA

fragment including the *rhtC* structural gene encoding the RhtC protein is ligated with a promoter sequence which functions efficiently in a bacterium belonging to the genus *Escherihia*. The Rt activity can be also enhanced
5 by substitution of the promoter sequence of the *rhtC* gene on a chromosome with a promoter sequence which functions efficiently in a bacterium belonging to the genus *Escherichia*.

Besides, the Rh actibity can be enhanced by, for
10 example, amplification of the copy number of the *rhtB* structural gene in a cell, or transformation of a bacterium belonging to the genus *Escherichia* with recombinant DNA in which a DNA fragment including the *rhtB* structural gene encoding RhtB protein is ligated
15 with a promoter sequence which functions efficiently in a bacterium belonging to the genus *Escherichia*. The Rh activity can be also enhanced by substitution of the promoter sequence of the *rhtB* gene on a chromosome with a promoter sequence which functions efficiently in a
20 bacterium belonging to the genus *Escherichia*.

The amplification of the copy number of the *rhtC* structural gene or *rhtB* structural gene in a cell can be performed by introduction of a multicopy vector which carries the *rhtC* structural gene or *rhtB* structural gene
25 into a cell of a bacterium belonging to the genus

Escherichia. Specifically, the copy number can be increased by introduction of a plasmid, a phage or a transposon (Berg, D.E. and Berg, C.M., Bio/Tecnol., 1, 417 (1983)) which carries the *rhtC* structural gene or
5 *rhtB* structural gene into a cell of a bacterium belonging to the genus *Escherichia*.

The multicopy vector is exemplified by plasmid vectors such as pBR322, pMW118, pUC19 or the like, and phage vectors such as λ 1059, λ BF101, M13mp9 or the like.
10 The transposon is exemplified by Mu, Tn10, Tn5 or the like.

The introduction of a DNA into a bacterium belonging to the genus *Escherichia* can be performed, for example, by a method of D.A M.Morrison (Methods in Enzymology, 68, 326 (1979)) or a method in which recipient bacterial
15 cell are treated with calcium chloride to increase permeability of DNA (Mandel, M. And Higa, A., J. Mol. Biol., 53, 159, (1970)) and the like.

If the Rt activity, or the Rt activity and the Rh
20 activity is enhanced in an amino acid-producing bacterium belonging to the genus *Escherichia* as described above, a produced amount of the amino acid can be increased. As the bacterium belonging to the genus *Escherichia* which is to be the Rt activity, or the Rt
25 activity and the Rh activity is enhanced, strains which

have abilities to produce desired amino acids are used. Besides, an ability to produce an amino acid may be imparted to a bacterium in which the Rt activity, of the Rt activity and Rh activity is enhanced.

5 On the basis of the rhtC DNA fragment amplification the new strains *E. coli* MG442/pRhtC producing homoserine; *E. coli* MG442/pVIC40,pRhtC producing threonine; *E. coli* NZ10/pRhtBC and *E. coli* NZ10/pRhtB, pRhtC producing homoserine, valine and leucine were
10 obtained which accumulate the amino acids in a higher amount than those containing no amplified rhtC DNA fragment.

 The new strains have been deposited (according to international deposition based on Budapest Treaty) in
15 the All-Russian Collection for Industrial Microorganisms (VKPM). The strain *E. coli* MG442/pRhtC has been deposited as an accession number of VKPM B-7700; the strain *E. coli* MG442/pVIC40,pRhtC has been deposited as an accession number of VKPM B-7680; the strain *E. coli*
20 NZ10/pRhtB, pRhtC has been deposited as an accession number of VKPM B-7681, and the strain *E. coli* NZ10/pRhtBC has been deposited as an accession number of VKPM B-7682.

 The strain *E. coli* MG442/pRhtC (VKPM B-7700)
25 exhibits the following cultural-morphological and

biochemical features.

Cytomorphology

Gram-negative weakly-motile rods having rounded
5 ends. Longitudinal size, 1.5 to 2 μm .

Cultural features

Beef-extract agar:

After 24 hours of growth at 37° C. produces round
10 whitish semitransparent colonies 1.0 to 3 mm in diameter,
featuring a smooth surface, regular or slightly wavy
edges, the centre is slightly
raised, homogeneous structure, pastelike consistency,
readily emulsifiable.

15

Luria's agar:

After a 24-hour growth at 37° C. develops whitish
semitranslucent colonies 1.5 to 2.5 mm in diameter
having a smooth surface, homogeneous structure,
20 pastelike consistency, readily emulsifiable.

Minimal agar-doped medium M9:

After 40 to 48 hours of growth at 37°C forms
colonies 0.5 to 1.5 mm in diameter, which are coloured
25 greyish-white, semitransparent, slightly convex, with a

lustrous surface.

Growth in a beaf-extract broth:

After a 24-hour growth at 37° C exhibits strong
5 uniform cloudiness, has a characteristic odour.

Physiological and biochemical features.

Grows upon thrust inoculation in a beef-extract agar:

Exhibits good growth throughout the inoculated
10 area. The microorganism proves to be a facultative
anaerobe.

It does not liquefy gelatin.

Features a good growth on milk, accompanied by milk
coagulation.

15 Does not produce indole.

Temperature conditions: Grows on beaf-extract broth at
20-42°C, an optimum temperature lying within 33-37 °C.

pH value of culture medium: Grows on liquid media
having the pH value from 6 to 8, an optimum value being
20 7.2.

Carbon sources: Exhibits good growth on glucose,
fructose, lactose, mannose, galactose, xylose, glycerol,
mannitol to produce an acid and gas.

Nitrogen sources: Assimilates nitrogen in the form of
25 ammonium, nitric acid salts, as well as from some

organic compounds.

Resistant to ampicillin.

L-isoleucine is used as a growth factor. However, the strain can grow slowly without isoleucine.

- 5 Content of plasmids: The cells contain multicopy hybrid plasmid pRhtC ensuring resistance to ampicillin (100 mg/l) and carrying the rhtC gene responsible for the increased resistance to threonine (50 mg/ml).

66224 22399
10 The strain *E. coli* MG442/pVIC40, pRhtC (VKPM B-7680) has the same cultural-morphological and biochemical features as the strain VKPM B-7700 except for in addition to pRhtC, it contains a multicopy hybrid plasmid pVIC40 ensuring resistance to streptomycin (100 mg/l) and carrying the genes of the threonine operon.

- 15 The strain *E. coli* strain *E. coli* NZ10/pRhtB, pRhtC (VKPM B-7681) has the same cultural-morphological and biochemical features as the strain VKPM B-7700 except for L-threonine (0.1 - 5 mg/ml) is used as a growth factor instead of L-isoleucine. Besides, it
20 contains a multicopy hybride plasmid pRhtB ensuring resistance to kanamycin (50 mg/l) and carrying the rhtB gene which confers resistance to homoserine (10 mg/ml)

The strain *E. coli* strain *E. coli* NZ10/pRhtBC, (VKPM B-7682) has the same cultural-morphological and
25 biochemical features as the strain VKPM B-7681 except

for it contains a multicopy hybride plasmid pRhtBC
ensuring resistance to ampicillin (100 mg/l) and
carrying both the rhtB and rhtC genes which confer
resistance to L-homoserine (10 mg/ml) and L-threonine
5 (50mg/ml).

<3> Method for producing an amino acid

An amino acid can be efficiently produced by
cultivating the bacterium in which the Rt activity, or
10 the Rt activity and Rh activity is enhanced by
amplifying a copy number of the *rhtC* gene, or *rhtC* gene
and *rhtB* gene as describe above, and which has an
ability to produce the amino acid, in a culture medium,
producing and accumulating the amino acid in the medium,
15 and recovering the amino acid from the medium. The amino
acid is exemplified preferably by L-homoserine, L-
threonine and branched chain amino acids. The branched
chain amino acids may be exemplified by L-valine, L-
leucine and L-isoleucine, and preferably exemplified by
20 L-valine, L-leucine.

In the method of present invention, the cultivation
of the bacterium belonging to the genus *Escherichia*, the
collection and purification of amino acids from the
liquid medium may be performed in a manner similar to
25 those of the conventional method for producing an amino

acid by fermentation using a bacterium. A medium used in cultivation may be either a synthetic medium or a natural medium, so long as the medium includes a carbon and a nitrogen source and minerals and, if necessary, nutrients which the bacterium used requires for growth in appropriate amount. The carbon source may include various carbohydrates such as glucose and sucrose, and various organic acids. Depending on assimilatory ability of the used bacterium. Alcohol including ethanol and glycerol may be used. As the nitrogen source, ammonia, various ammonium salts as ammonium sulfate, other nitrogen compounds such as amines, a natural nitrogen source such as peptone, soybean hydrolyzate and digested fermentative microbe are used. As minerals, monopotassium phosphate, magnesium sulfate, sodium chloride, ferrous sulfate, manganese sulfate, calcium carbonate are used.

The cultivation is preferably culture under an aerobic condition such as a shaking, and an aeration and stirring culture. The temperature of culture is usually 20 to 40°C, preferably 30 to 38°C. The pH of the culture is usually between 5 and 9, preferably between 6.5 and 7.2. the pH of the culture can be adjusted with ammonia, calcium carbonate, various acids, various bases, and buffers. Usually, a 1 to 3-day cultivation leads to the

accumulation of the target amino acid in the medium.

Recovering the amino acid can be performed by removing solids such as cells from the medium by centrifugation or membrane filtration after cultivation, and then collecting and purifying the target amino acid by ion exchange, concentration and crystalline fraction methods and the like.

Brief Explanation of Drawings

10

Fig. 1 shows cloning and identification of *rhtB* and *rhtC* genes,

Fig. 2 shows structure of the plasmid pRhtB harboring *rhtB* gene,

15

Fig. 3 shows structure of the plasmid pRhtC harboring *rhtC* gene, and

Fig. 4 shows structure of the plasmid pRhtBC harboring *rhtB* gene and *rhtC* gene.

20

Best Mode for Carrying Out the Invention

The present invention will be more concretely explained below with reference to Examples. In the Examples, an amino acid is of L-configuration unless otherwise noted.

25

Example 1: Obtaining of the rhtB and rhtC DNA fragmentsStep 1. Cloning of genes involving resistance homoserine
5 and threonine into mini-Mu phagemid

The genes involving resistance homoserine and threonine were cloned *in vivo* using mini-Mu d5005 phagemid (Groisman, E.A., et al., J. Bacteriol., 168, 357-364 (1986)). MuCts62 lysogen of the strain MG442
10 (Guayatiner et al., Genetika (in Russian), 14, 947-956 (1978)) was used as a donor. Freshly prepared lysates were used to infect a MuCts lysogenic derivative of a strain VKPM B-513 (Hfr K10 metB). The cells were plated on M9 glucose minimal medium with methionine (50 µg/ml),
15 kanamycin (40 µg/ml) and homoserine (10 µg/ml). Colonies which appeared after 48 hr were picked and isolated. Plasmid DNA was isolated and used to transform the strain VKPM B-513 by standard techniques. Transformants were selected on L-broth agar plates with kanamycin as
20 above. Plasmid DNA was isolated from those which were resistance to homoserine, and analyzed by restriction mapping of the structure of the inserted fragments. It appeared that two types of inserts belonging to different chromosome regions had been cloned from the

donor. Thus, at least two different genes that in multicopy impart resistance to homoserine exist in *E. coli*. One of the two types of inserts is the *rhtA* gene which has already reported (ABSTRACT of 17th International Congress of Biochemistry and Molecular Biology in conjugation with 1997 Annual Meeting of the American Society for Biochemistry and Molecular Biology, San Francisco, California August 24-29, 1997). Among the other of the two types of inserts, a MluI-MluI fragment of 0.8 kb imparts only the resistance to homoserine (Fig. 1).

Step 2: Identification of *rhtB* gene and *rhtC* gene

The insert fragment was sequenced by the dideoxy chain termination method of Sanger. Both DNA strands were sequenced in their entirety and all junctions were overlapped. The sequencing showed that the insert fragment included fl38 (nucleotide numbers 61543 to 61959 of GenBank accession number M87049) which was a known but function-unknown ORF (open reading frame) present at 86 min of *E. coli* chromosome and about 350 bp of an upstream region thereof (downstream region in the sequence of M87049). The fl38 which had only 160 nucleotides in the 5'-flanking region could not impart the resistance to homoserine. No termination codon is

present upstream the ORF fl38 between 62160 and 61950 nucleotides of M87049. Furthermore, one ATG following a sequence predicted as a ribosome binding site is present in the sequence. The larger ORF (nucleotide numbers
 5 62160 to 61546) is designated as *rhtB* gene. The RhtB protein deduced from the gene has a region which is highly hydrophobic and contains possible transmembrane segments.

As described below, the plasmid containing this gene
 10 conferred upon cells only the resistance to high concentrations of homoserine. Since the initial *Sac*II-*Sac*II DNA fragment contained the second unidentified ORF, 0128, the gene was subcloned and tested for its ability to confer resistance to homoserine and threonine. It
 15 proved that the plasmid containing 0128 (*Cla*I-*Eco*47III fragment) conferred resistance to 50 mg/ml threonine (Fig. 1). The subcloned fragment was sequenced and found to contain additional nucleotide (G) in the position between 61213 and 61214 nucleotides of M87049. The
 20 nucleotide addition to the sequence eliminated a frame shift and enlarged the ORF into 5'-flanking region up to 60860 nucleotide. This new gene was designated as *rhtC*. Both genes, *rhtB* and *rhtC*, were found to be homologous to transporter involved in lysine export of
 25 *Corynebacterium glutamicum*.

Example 2: The effect of *rhtB* and *rhtC* genes
amplification on homoserine production.

<1> Construction of the L-homoserine-producing strain *E.*

5 *coli* NZ10/pAL4, pRhtB and homoserine production

The *rhtB* gene was inserted to a plasmid pUK21
(Vieira, J. And Messing, J., Gene, 100, 189-194 (1991)),
to obtain pRhtB (Fig. 2).

Strain NZ10 of *E. coli* was transformed by a plasmid
10 pAL4 which was a pBR322 vector into which the *thrA* gene
coding for aspartokinase-homoserine dehydrogenase I was
inserted, to obtain the strains NZ10/pAL4. The strain
NZ10 is a *leuB*⁺-reverted mutant *thrB*⁻ obtained from the
E. coli strain C600 (*thrB*, *leuB*) (Appleyard R.K.,
15 Genetics, 39, 440-452, 1954).

The strain NZ10/pAL4 was transformed with pUK21 or
pRhtB to obtain strains NZ10/pAL4,pUK21 and NZ10/pAL4,
pRhtB.

The thus obtained transformants were each cultivated
20 at 37°C for 18 hours in a nutrient broth with 50 mg/l
kanamycin and 100 mg/l ampicilin, and 0.3 ml of the
obtained culture was inoculate into 3 ml of a
fermentation medium having the following composition and
containing 50 mg/l kanamycin and 100 mg/l ampicilin, in
25 a 20 x 200 mm test tube, and cultivated at 37°C for 48

hours with a rotary shaker. After the cultivation, an accumulated amount of homoserine in the medium and an absorbance at 560 nm of the medium were determined by known methods.

5

[Fermentation medium composition (g/L)]

	Glucose	80
	(NH ₄) ₂ SO ₄	22
	K ₂ HPO ₄	2
10	NaCl	0.8
	MgSO ₄ •7H ₂ O	0.8
	FeSO ₄ •7H ₂ O	0.02
	MnSO ₄ •5H ₂ O	0.02
	Thiamine hydrochloride	0.2
15	Yeast Extract	1.0
	CaCO ₃	30
	(CaCO ₃ was separately sterilized)	

The results are shown in Table 1. As shown in Table 1, the strain NZ10/pAL4,pRhtB accumulated homoserine in a larger amount than the strain NZ10/pAL4,pUK21 in which the *rhtB* gene was not enhanced.

25

Table 1.

Strain	OD ₅₆₀	Accumulated amount of homoserine(g/L)
NZ10/pAL4,pUK21	14.3	3.3
NZ10/pAL4,pRhtB	15.6	6.4

<2> Construction of the homoserine-producing strain *E. coli* MG442/pRhtC and homoserine production

5 The *rhtC* gene was inserted to pUC21 vector (Vieira, J. And Messing, J., *Gene*, 100, 189-194 (1991)), to obtain pRhtC (Fig. 3).

The known *E. coli* strain MG442 which can produce threonine in an amount of not less than 3 g/L
 10 (Gusyatiner, et al., 1978, *Genetika* (in Russian), 14:947-956) was transformed by introducing pUC21 or pRhtC to obtain the strains MG442/pUC21 and MG442/pRhtC.

The thus obtained transformants were each cultivated at 37°C for 18 hours in a nutrient broth with 100 mg/ml
 15 ampicilin, and 0.3 ml of the obtained culture was inoculate into 3 ml of a fermentation medium describe above and containing 100 mg/ml ampicilin, in a 20 x 200 mm test tube, and cultivated at 37°C for 48 hours with a rotary shaker. After the cultivation, an accumulated
 20 amount of homoserine in the medium and an absorbance at 560 nm of the medium were determined by known methods. The results are shown in Table 2.

Table 2.

Strain	OD ₅₆₀	Accumulated amount of homoserine (g/L)
MG442/pUC21	9.7	<0.1
MG442/pRhtC	15.2	9.5

Example 3: The effect of rhtB and rhtC genes
amplification on threonine production.

5

<1> Construction of the threonine -producing strain *E. coli* VG442/pVIC40, pRhtB (VKPM B-7660) and threonine production

The strain MG442 was transformed by introducing a
10 known plasmid pVIC40 (U.S. Patent No. 5,175,107 (1992))
by an ordinary transformation method. Transformants were
selected on LB agar plates containing 0.1 mg/ml
streptomycin. Thus a novel strain MG442/pVIC40 was
obtained.

15 The strain MG442/pVIC40 was transformed with pUK21
or pRhtB to obtain strain MG442/pVIC40,pUK21 and
MG442/pVIC40,pRhtB.

The thus obtained transformants were each cultivated
at 37°C for 18 hours in a nutrient broth with 50 mg/l
20 kanamycin and 100 mg/l streptomycin, and 0.3 ml of the
obtained culture was inoculate into 3 ml of a

fermentation medium describe in Example 2 and containing 50 mg/l kanamycin and 100 mg/l streptomycin, in a 20 x 200 mm test tube, and cultivated at 37°C for 68 hours with a rotary shaker. After the cultivation, an accumulated amount of threonine in the medium and an absorbance at 560 nm of the medium were determined by known methods.

The results are shown in Table 3. As shown in Table 3, the strain MG442/pVIC40,pRhtB accumulated threonine in a larger amount than the strain MG442/pVIC40,pUK21 in which the *rhtB* gene was not enhanced.

Table 3.

Strain	OD ₅₆₀	Accumulated amount of threonine (g/L)
MG442/pVIC40,pUK21	16.3	12.9
MG442/pVIC40,pRhtB	15.2	16.3

<2> Construction of the threonine-producing strain *E. coli* VG442/pVIC40, pRhtC (VKPM B-7680) and threonine production

The strain MG442/pVIC40 was transformed with pRhtC and pUC21. Thus the transformants MG442/pVIC40,pRhtC and MG442/pVIC40, pUC21 were obtained. In the sane manner as describe above, MG442/pVIC40,pUC21 and

MG442/pVIC40,pRhtC were each cultivated at 37°C for 18 hours in a nutrient broth with 100 mg/l ampicilin and 100 mg/l streptomycin and 0.3 ml of the obtained culture was inoculate into 3 ml of a fermentation medium describe above and containing 100 mg/l ampicilin and 100 mg/l streptomycin, in a 20 x 200 mm test tube, and cultivated at 37°C for 46 hours with a rotary shaker. After the cultivation, an accumulated amount of threonine in the medium and an absorbance at 560 nm of the medium were determined by known methods.

The results are shown in Table 4. As shown in Table 4, the strain MG442/pVIC40,pRhtC accumulated threonine in a larger amount than the strain MG442/pVIC40,pUC21 in which the *rhtC* gene was not enhanced.

Table 4

Strain	OD ₅₆₀	Accumulated amount of threonine (g/L)
MG442/pVIC40, pUC21	17.4	4.9
MG442/pVIC40,pRhtC	15.1	10.2

Example 4: Concerted effect of *rhtB* gene and *rhtC* gene on amino acid production

The *Sac*II-*Sac*II DNA fragment containing both *rhtB* and *rhtC* genes was inserted to the pUC21. Thus the plasmid pRhtBC was obtained which harbors the *rhtB* gene

and *rhtC* gene (Fig. 4).

Then, the strain NZ10 was transformed with pUC21, pRhtB, pRhtC or pRhtBC, and the transformants NZ10/pUC21 (VKPM B-7685), NZ10/pRhtB (VKPM B-7683), NZ10/pRhtC
5 (VKPM B-7684), NZ10/pRhtB, pRhtC (VKPM B-7681) and NZ10/pRhtBC (VKPM B-7682) were thus obtained.

The transformants obtained above were cultivated as the same manner as describe above and accumulated amounts of various amino acids in the medium and an
10 absorbance at 540 nm of the medium were determined by known methods.

The result were shown in Table 5. It follows from the Table 5 that concerted effect of the pRhtB and pRhtC on producrion of homoserine, valine and leucine. These
15 results indicate that the *rhtB* and *rhtC* gene products may interact in cells.

Table 5.

Strain	OD ₅₆₀	Homoserine (g/L)	Valine (g/L)	Leucine (g/L)
NZ10/pUC21	18.7	0.6	0.22	0.16
NZ10/pRhtB	19.6	2.3	0.21	0.14
NZ10/pRhtC	20.1	0.7	0.2	0.15
NZ10/pRhtBC	21.8	4.2	0.34	0.44
NZ10/pRhtB,pRhtC	19.2	4.4	0.35	0.45

Example 5: Effect of rhtB gene and rhtC gene on
resistance to amino acids

5

As describe above, the plasmids harboring the *rhtB* and *rhtC* have positive effect on some amino acid accumulation in culture broth by different strains. It proved that the pattern of accumulated amino acid was dependent on the strain genotype. The homology of the *rhtB* and *rhtC* genes products with the lysine transporter LysE of *Corynebacterium glutamicum* (Vrljic, M., Sahm, H. and Eggeling, L. (1996) *Mol. Microbiol.* 22, 815-826.) indicates the analogues function for these proteins.

15 Therefore, the effect of the pRhtB and pRhtC plasmids on susceptibility of the strain N99 which is a streptomycin-resistant (Str^R) mutant of the known strain W3350 (VKPM B-1557) to some amino acids and amino acid

analogues was tested. Overnight broth cultures (10^9 cfu/ml) of the strains N99/pUC21, N99pUK21, N99/pRhtB and N99/pRhtC were diluted 1:100 in M9 minimal medium and grown for 5 h in the same medium. Then the log phase cultures thus obtained were diluted and about 10^4 viable cells were applied to well-dried test plates with M9 agar (2%) containing doubling increments of amino acids or analogues. Thus the minimum inhibitory concentration (MIC) of these compounds were examined.

10 The result are shown in Table 6. It follows from the Table 6 that multiple copies of *rhtB* besides homoserine conferred increased resistance to α -amino- β -hydroxyvaleric-acid (AHVA) and S-(2-aminoethyl)-L-cysteine (AEC), and 4-aza-DL-leucine; and multiple
15 copies of *rhtC* gene besides threonine increased resistance to valine, histidine, and AHVA. This results indicates that every of the presumed transporters, *RhtB* and *RhtC*, have specificity to several substrates (amino acids), or may shown non-specific effects as a result of
20 amplification.

Table 6.

Substrate	MIC ($\mu\text{g/ml}$)		
	N99/pUC21*	N99/pRhtB	N99/pRhtC
L-homoserine	1000	20000	1000
L-threonine	30000	40000	80000
L-valine	0.5	0.5	2.0
L-histidine	5000	5000	40000
AHVA	100	2000	15000
AEC	5	20	5
4-aza-DL-leucine	50	100	50
O-methyl-L-threonine	20	20	20

*: The same data were obtain with N99/pUK21.

SEQUENCE LISTING

<110> Livshits, Vitaliy Arkadyevich
 Zakataeva, Natalia Pavlovna
 Aleshin, Vladimir Veniaminovich
 Belareva, Alla Valentinovna
 Tokhmakova, Irina Lyvovna

<120> NOVEL GENE AND METHOD FOR PRODUCING L-AMINO ACIDS

<130> OP851

<141> 1999-12-

<150> RU-98123511

<151> 1998-12-23

<160> 4

<170> PatentIn Ver. 2.0

<210> 1

<211> 1231

<212> DNA

<213> Escherichia coli

<220>

<221> CDS

<222> (557)..(1171)

<400> 1

```

agaaataatg tggagatcgc accgcccatac gaatgtgccg gtatatagcg tttagccac 60
ggaccgggct gaacctctg ctgccagaat gccgccagat catcaacata atcattaaag 120
cgattaacat gcccgagatg cggatcggt aacaggcgac cggaacgtcc ctgcccgcga 180
tggtcgatga ttaagacatc aaaccccaaa tggaacaggt cataggccag ttccgcata 240
tttagctagc tctcaatacg ccccgggcag atgactacca cccgggtcatg gtgctgtgcg 300
cgaaaacgga caaagcgcac cggaatgtca tccacaccag taaactctgc ttcacacgc 360
tgacgccaga aatcagtcag cggtcctatg gtaaaagcag caaacgcgtt ttctctgtt 420
tcccagtcct tttgtgctg aaacatcggg taatctgcct cttaaaccac gtaaaatcgt 480
tttttttagc gtgcctgaca caacgtgcg acagtagcgt attgtggcac aaaaatagac 540
acaccgggag ttcatac atg acc tta gaa tgg tgg ttt gcc tac ctg ctg aca 592
                                Met Thr Leu Glu Trp Trp Phe Ala Tyr Leu Leu Thr
                                1           5           10
tcg atc att tta acg ctg tgc cca ggc tct ggt gca atc aac act atg 640
Ser Ile Ile Leu Thr Leu Ser Pro Gly Ser Gly Ala Ile Asn Thr Met

```

<400> 2
Met Thr Leu Glu Trp Trp Phe Ala Tyr Leu Leu Thr Ser Ile Ile Leu
1 5 10 15

tgc ggc gta atg gtt tgg gct ggg att gcg ctg ctt ggc ctg cat ttg 372
 Cys Gly Val Met Val Trp Ala Gly Ile Ala Leu Leu Gly Leu His Leu
 50 55 60
 att atc gaa aaa atg gcc tgg ctg cat acg ctg att atg gtg ggc ggt 420
 Ile Ile Glu Lys Met Ala Trp Leu His Thr Leu Ile Met Val Gly Gly
 65 70 75
 ggc ctg tat ctc tgc tgg atg ggt tac cag atg cta cgt ggt gca ctg 468
 Gly Leu Tyr Leu Cys Trp Met Gly Tyr Gln Met Leu Arg Gly Ala Leu
 80 85 90
 aaa aaa gag gcg gtt tct gca cct gcg cca cag gtc gag ctg gcg aaa 516
 Lys Lys Glu Ala Val Ser Ala Pro Ala Pro Gln Val Glu Leu Ala Lys
 95 100 105 110
 agt ggg cgc agt ttc ctg aaa ggt tta ctg acc aat ctc gct aat ccg 564
 Ser Gly Arg Ser Phe Leu Lys Gly Leu Leu Thr Asn Leu Ala Asn Pro
 115 120 125
 aaa gcg att atc tac ttt ggc tcg gtg ttc tca ttg ttt gtc ggt gat 612
 Lys Ala Ile Ile Tyr Phe Gly Ser Val Phe Ser Leu Phe Val Gly Asp
 130 135 140
 aac gtt ggc act acc gcg cgc tgg ggc att ttt gcg ctg atc att gtc 660
 Asn Val Gly Thr Thr Ala Arg Trp Gly Ile Phe Ala Leu Ile Ile Val
 145 150 155
 gaa acg ctg gcg tgg ttt acc gtc gtt gcc agc ctg ttt gcc ctg ccg 708
 Glu Thr Leu Ala Trp Phe Thr Val Val Ala Ser Leu Phe Ala Leu Pro
 160 165 170
 caa atg cgc cgt ggt tat caa cgt ctg gcg aag tgg att gat ggt ttt 756
 Gln Met Arg Arg Gly Tyr Gln Arg Leu Ala Lys Trp Ile Asp Gly Phe
 175 180 185 190
 gcc ggg gcg tta ttt gcc gga ttt ggc att cat ttg att att tcg cgg 804
 Ala Gly Ala Leu Phe Ala Gly Phe Gly Ile His Leu Ile Ile Ser Arg
 195 200 205
 tgatgccaga cgcgtcttca gagtaagtcg gataag 840

<210> 4

<211> 206

<212> PRT

<213> Escherichia coli

<400> 4

Met Leu Met Leu Phe Leu Thr Val Ala Met Val His Ile Val Ala Leu
 1 5 10 15
 Met Ser Pro Gly Pro Asp Phe Phe Phe Val Ser Gln Thr Ala Val Ser
 20 25 30
 Arg Ser Arg Lys Glu Ala Met Met Gly Val Leu Gly Ile Thr Cys Gly
 35 40 45
 Val Met Val Trp Ala Gly Ile Ala Leu Leu Gly Leu His Leu Ile Ile
 50 55 60
 Glu Lys Met Ala Trp Leu His Thr Leu Ile Met Val Gly Gly Gly Leu

What is claimed is:

1. A bacterium belonging to the genus *Escherichia*, wherein L-threonine resistance of said bacterium is enhanced by enhancing an activity of protein as defined in the following (A) or (B) in a cell of said bacterium:

(A) a protein which comprises the amino acid sequence shown in SEQ ID NO: 4 in Sequence Listing; or

(B) a protein which comprises the amino acid sequence including deletion, substitution, insertion or addition of one or several amino acids in the amino acid sequence shown in SEQ ID NO: 4 in Sequence Listing, and which has an activity of making a bacterium having the protein L-threonine-resistant.

2. The bacterium according to claim 1, wherein L-homoserine resistance of said bacterium is further enhanced by enhancing an activity of protein as defined in the following (C) or (D) in a cell of said bacterium:

(C) a protein which comprises the amino acid sequence shown in SEQ ID NO: 2 in Sequence Listing; or

(D) a protein which comprises the amino acid sequence including deletion, substitution, insertion or addition of one or several amino acids in the amino acid sequence shown in SEQ ID NO: 2 in Sequence Listing, and which has an activity of making a bacterium having the protein L-homoserine-resistant.

3. The bacterium according to claim 1 or 2, wherein said activity of protein as defined in (A) or (B) is enhanced by transformation of said bacterium with DNA coding for the protein as defined in (A) or (B).

4. The bacterium according to claim 2, wherein said activity of protein as defined in (C) or (D) is enhanced by transformation of said bacterium with DNA coding for the protein as defined in (C) or (D).

5. A method for producing an amino acid, comprising the steps of:

cultivating the bacterium as defined in any one of claims 1 to 4, which has an ability to produce the amino acid, in a culture medium, to produce and accumulate the amino acid in the medium, and

recovering the amino acid from the medium.

6. The method according to claim 5, wherein said amino acid is selected from the group consisting of L-homoserine, L-threonine and branched chain amino acids.

7. The method according to claim 6, said branched chain amino acids is L-valine or L-leucine.

8. A DNA which encode a protein defined in the following (A) or (B):

(A) a protein which has the amino acid sequence of SEQ ID NO: 4;

(B) a protein which has the amino acid sequence of SEQ ID NO: 4 including substitution, deletion, insertion, addition, or inversion of one or several amino acids,

9. The DNA of claim 8 which is a DNA defined in the following (a) or (b):

(b) a DNA which is hybridizable with a nucleotide sequence of nucleotide numbers 187 to 804 in SEQ ID NO: 3 or a probe prepared from the nucleotide sequence under a stringent condition, and encodes a protein having an activity of making a bacterium having the protein L-threonine-resistant.

10. The DNA of claim 9 wherein the stringent condition is a condition in which washing is performed at 60 °C, and at a salt concentration corresponding to 1 x SSC and 0.1 % SDS.

Abstract

A bacterium which has an ability to produce an amino acid and in which *rhtC* gene coding for a protein having an activity of making a bacterium having the protein L-threonine-resistant is enhanced, preferably, in which *rhtB* gene coding for a protein having an activity of making a bacterium having the protein L-homoserine-resistant is further enhanced, is cultivated in a culture medium to produce and accumulate the amino acid in the medium, and the amino acid is recovered from the medium.




Fig. 1




Fig. 2




Fig. 3




Fig. 4

Declaration, Power Of Attorney and Petition

Page 1 of 3

WE (I) the undersigned inventor(s), hereby declare(s) that:

My residence, post office address and citizenship are as stated below next to my name,

We (I) believe that we are (I am) the original, first, and joint (sole) inventor(s) of the subject matter which is claimed and for which a patent is sought on the invention entitled

NOVEL GENE AND METHOD FOR PRODUCING L-AMINO ACIDS

the specification of which

☒ is attached hereto.

☐ was filed on _____ as

Application Serial No. _____

and amended on _____.

☐ was filed as PCT international application

Number _____

on _____,

and was amended under PCT Article 19

on _____ (if applicable).

We (I) hereby state that we (I) have reviewed and understand the contents of the above-identified specification, including the claims, as amended by any amendment referred to above.

We (I) acknowledge the duty to disclose information known to be material to the patentability of this application as defined in Section 1.56 of Title 37 Code of Federal Regulations.

We (I) hereby claim foreign priority benefits under 35 U.S.C. § 119(a)-(d) or § 365(b) of any foreign application(s) for patent or inventor's certificate, or § 365(a) of any PCT International application which designated at least one country other than the United States, listed below and have also identified below, by checking the box, any foreign application for patent or inventor's certificate, or PCT International application having a filing date before that of the application on which priority is claimed. Prior Foreign Application(s)

Application No.	Country	Day/Month/Year	Priority Claimed
98123511	Russia	23/12/1998	<input checked="" type="checkbox"/> Yes <input type="checkbox"/> No
_____	_____	_____	<input type="checkbox"/> Yes <input type="checkbox"/> No
_____	_____	_____	<input type="checkbox"/> Yes <input type="checkbox"/> No
_____	_____	_____	<input type="checkbox"/> Yes <input type="checkbox"/> No

We (I) hereby claim the benefit under Title 35, United States Code, § 119(e) of any United States provisional application(s) listed below.

_____	_____
(Application Number)	(Filing Date)
_____	_____
(Application Number)	(Filing Date)

We (I) hereby claim the benefit under 35 U.S.C. § 120 of any United States application(s), or § 365(c) of any PCT International application designating the United States, listed below and, insofar as the subject matter of each of the claims of this application is not disclosed in the prior United States or PCT International application in the manner provided by the first paragraph of 35 U.S.C. § 112, I acknowledge the duty to disclose information which is material to patentability as defined in 37 CFR § 1.56 which became available between the filing date of the prior application and the national or PCT International filing date of this application.

Application Serial No.	Filing Date	Status (pending, patented, abandoned)
_____	_____	_____
_____	_____	_____
_____	_____	_____


And we (I) hereby appoint: Norman F. Oblon, Registration Number 24,618; Marvin J. Spivak, Registration Number 24,913; C. Irvin McClelland, Registration Number 21,124; Gregory J. Maier, Registration Number 25,599; Arthur I. Neustadt, Registration Number 24,854; Richard D. Kelly, Registration Number 27,757; James D. Hamilton, Registration Number 28,421; Eckhard H. Kuesters, Registration Number 28,870; Robert T. Pous, Registration Number 29,099; Charles L. Gholz, Registration Number 26,395; Vincent J. Sunderdick, Registration Number 29,004; William E. Beaumont, Registration Number 30,996; Steven B. Kelber, Registration Number 30,073; Robert F. Gnuse, Registration Number 27,295; Jean-Paul Lavalleye, Registration Number 31,451; Timothy R. Schwartz, Registration Number 32,171; Stephen G. Baxter, Registration Number 32,884; Martin M. Zoltick, Registration Number 35,745; Robert W. Hahl, Registration Number 33,893; Richard L. Treanor, Registration Number 36,379; Steven P. Weihrouch, Registration Number 32,829; John T. Goolkasian, Registration Number 26,142; Marc R. Labgold, Registration Number 34,651; William J. Healey, Registration Number 36,160; and Richard L. Chinn, Registration Number 34,305; our (my) attorneys, with full powers of substitution and revocation, to prosecute this application and to transact all business in the Patent Office connected therewith; and we (I) hereby request that all correspondence regarding this application be sent to the firm of OBLON, SPIVAK, MCCLELLAND, MAIER & NEUSTADT, P.C., whose Post Office Address is: Fourth Floor, 1755 Jefferson Davis Highway, Arlington, Virginia 22202.

We (I) declare that all statements made herein of our (my) own knowledge are true and that all statements made on information and belief are believed to be true; and further that these statements were made with the knowledge that willful false statements and the like so made are punishable by fine or imprisonment, or both, under Section 1001 of Title 18 of the United States Code and that such willful false statements may jeopardize the validity of the application or any patent issuing thereon.

Vitaliy Arkadyevich LIVSHITS

NAME OF FIRST SOLE INVENTOR

Residence: Moscow, Russia



Signature of Inventor

Citizen of: Russia

Post Office Address: c/o Institute of Genetics and Selection of Industrial Microorganisms (GNIIGenetika), 1, Dorozhny pr., 1, 113545, Moscow, Russia

November 25, 1999

Date

Natalia Pavlovna ZAKATAEVA

NAME OF SECOND JOINT INVENTOR

Zakataeva

Signature of Inventor

November 25, 1999

Date

Vladimir Veniaminovich ALESHIN

NAME OF THIRD JOINT INVENTOR

Aleshin

Signature of Inventor

November 25, 1999

Date

Alla Valentinovna BELAREVA

NAME OF FOURTH JOINT INVENTOR

Belareva

Signature of Inventor

November 25, 1999

Date

Irina Lyvovna TOKHMAKOVA

NAME OF FIFTH JOINT INVENTOR

Tokhmakova

Signature of Inventor

November 25, 1999

Date

Residence: Moscow, Russia

Citizen of: Russia

Post Office Address: c/o Institute of Genetics and
Selection of Industrial Microorganisms (GNIIGenetika),
1, Dorozhny pr., 1, 113545, Moscow, Russia

Residence: Moscow, Russia

Citizen of: Russia

Post Office Address: c/o Institute of Genetics and
Selection of Industrial Microorganisms (GNIIGenetika),
1, Dorozhny pr., 1, 113545, Moscow, Russia

Residence: Moscow, Russia

Citizen of: Russia

Post Office Address: c/o Institute of Genetics and
Selection of Industrial Microorganisms (GNIIGenetika),
1, Dorozhny pr., 1, 113545, Moscow, Russia

Residence: Moscow, Russia

Citizen of: Russia

Post Office Address: c/o Institute of Genetics and
Selection of Industrial Microorganisms (GNIIGenetika),
1, Dorozhny pr., 1, 113545, Moscow, Russia